

# Hand Span Influences Optimal Grip Span in Boys and Girls Aged 6 to 12 Years

Vanesa España-Romero, BS, Enrique G. Artero, BS, Alba M. Santaliestra-Pasias, BS, Angel Gutierrez, MD, PhD, Manuel J. Castillo, MD, PhD, Jonatan R. Ruiz, PhD

**Purpose** The first aim was to determine whether there is an optimal grip span for determining the maximum hand grip strength in boys and girls aged 6 to 12 years and whether the optimal grip span was related to hand span. If so, the second aim was to derive a mathematical equation relating hand span and optimal grip span.

**Methods** A total of 123 boys ( $9 \text{ y} \pm 2$ ) and 70 girls ( $8 \text{ y} \pm 2$ ) were evaluated. Each hand was randomly tested on 10 occasions using 5 different grip spans, allowing a 1-minute rest between attempts. The hand span was measured from the tip of the thumb to the tip of the little finger with the hand opened widely.

**Results** An optimal grip span to determine maximum hand grip strength was identified for both genders. Hand span and optimal grip span showed a significant linear association in the studied children. The equation relating grip span as a function of hand span in boys is formulated as  $y = x/4 + 0.44$  and in girls as  $y = 0.3x - 0.52$ , where  $x$  is the hand span (maximal width between first and fifth fingers) and  $y$  is the optimal grip span.

**Conclusions** The results suggest that there is an optimal grip span to which the dynamometer should be adjusted when measuring hand grip strength in children. The optimal grip span was influenced by hand span in both genders. (*J Hand Surg* 2008;33A:378–384. Copyright © 2008 by the American Society for Surgery of the Hand.)

**Key words** Children, dynamometry, hand grip strength, reliability, standardization.

**H**AND GRIP STRENGTH is a widely used test in experimental and epidemiologic studies in children.<sup>1–3</sup> The measure of hand grip strength is influenced by several factors, including age; gender; different angle of shoulder, elbow, forearm,

and wrist; and posture.<sup>4–9</sup> Other important factors affecting hand grip strength are grip span<sup>8,10,11</sup> and hand span.<sup>12–15</sup> Härkönen et al<sup>11</sup> showed that hand grip strength varied with hand grip position and was slightly affected by hand span. We have shown that there is an optimal grip span at which the maximum hand grip strength is obtained in both teenagers<sup>12</sup> and in adults.<sup>13</sup> Therefore, one would expect that children may need a different optimal grip span when measuring hand grip strength compared with older populations. Studies examining which position on the grip dynamometer will result in maximum hand grip strength in children are lacking.

The first aim of the current study was to determine if there is an optimal grip span for determining the maximum hand grip strength in boys and girls aged 6 to 12 years and if that grip span is related to hand span. If so, the second aim was to derive a mathematical equation relating hand span and optimal grip span.

## MATERIALS AND METHODS

### Subjects

A total of 123 boys ( $9 \text{ y} \pm 2$ ), and 70 girls ( $8 \text{ y} \pm 2$ ) participated in the study. A comprehensive verbal description of the nature and purpose of the study was given to both the children and their teachers. One parent or legal guardian provided written informed consent, and all children gave verbal assent. All the children included in the current study were in good health and free of any lesion or

*From the Department of Physiology, School of Medicine, University of Granada, Granada, Spain; the Unit for Preventive Nutrition, Department of Biosciences and Nutrition at NOVUM, Karolinska Institute, Huddinge, Sweden; and the E. U. Ciencias de la Salud, University of Zaragoza, Zaragoza, Spain.*

Received for publication October 10, 2007; accepted in revised form November 19, 2007.

This study was supported by the IDEFICS study, with the financial support of the European Community Sixth RTD Framework Programme (contract FOOD-CT-2006-016181-2); the ALPHA study, with the financial support of the Public Health Executive Agency, DG Sanco, Health Information Strand (ref. 2006120); Ministerio de Educación y Ciencia de España (AP2003-2128 and AP2005-4358); and the Margit and Folke Pehrzon Foundation. The content of this article reflects only the authors' views, and the European community is not liable for any use that may be made of the information contained herein.

**Corresponding author:** Vanesa España-Romero, BS, Department of Physiology, School of Medicine, University of Granada, 18071 Granada, Spain; e-mail: [vanesaespana05@yahoo.es](mailto:vanesaespana05@yahoo.es).

0363-5023/08/33A03-0014\$34.00/0  
doi:10.1016/j.jhsa.2007.11.013




FIGURE 1: Measurement of hand span (0.5-cm precision).

impairments in the upper limbs. The children were encouraged to do their best when performing the tests. The study protocol was performed in accordance with the ethical standards established in the 1961 Declaration of Helsinki (as revised in Hong Kong in 1989 and in Edinburgh, Scotland, in 2000) and was approved by the Review Committee for Research Involving Human Subjects at the University of Granada.

## Methods

**Measurement of hand span:** Hand span was measured in both hands from the tip of the thumb to the tip of the small finger with the hand opened as wide as possible (Fig. 1). The accuracy of the measure was 0.5 cm. The results of hand span were therefore rounded to the nearest whole centimeter.

**Measurement of grip span:** Grip span was measured from the external base of the grip to the mark placed in the central strip of the dynamometer (see arrow in Fig. 2).

**Measurement of hand grip strength:** Hand grip strength was measured using a digital hand dynamometer (T.K.K. 5401 Grip-D; Takey, Tokyo, Japan), and the scores were recorded in kilograms. The reported precision of the dynamometer was 0.1 kg.



FIGURE 2: Device for measurement of grip span (0.1-cm precision).




FIGURE 3: Association relating hand grip strength and grip span in 1 of the cases (boy, 7 years old, right hand). The maximum of the second-degree polynomial regression equation relating hand grip strength and grip span,  $f'(x)$ , was the optimal grip span for each hand of each individual.

A detailed description of the test can be found elsewhere.<sup>12,13</sup> In brief, the children were standing during the entire test with the arm at the side with the shoulder slightly abducted ( $\sim 10^\circ$ ), the elbow extended, and the forearm and wrist in neutral position. Each child performed the test twice (alternately with both hands) using different grip spans in random order and allowing a 1-minute rest between measures. For each measure, the hand to be tested first was chosen randomly. The grip spans used were 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, and 6.5 cm. If the hand span was bigger than 17 cm, then the lowest grip span was rejected; if the hand span was smaller than 17 cm, then the highest grip span was rejected. Furthermore, if the hand span was smaller than 15 cm, then the grip spans used ranged between 3.5 and 5.5 cm. For each hand, the best result for each grip span was retained.

The grip span equivalence for the different positions in the Jamar hand dynamometer (Fit System Inc., Calgary, Canada) are position 1, 3.5 cm; position 2, 4.8 cm; position 3, 6.0; position 4, 7.3 cm; and position 5, 8.6 cm.

**Determination of the optimal grip span:** The optimal grip span is the grip span at which the maximum hand grip strength is obtained. To determine the individual optimal grip span for each hand of each child, we first established the kind of association relating grip span and hand grip strength (ie, the results of hand grip strength obtained at the different grip spans). For that purpose, the statistical software SPSS ver. 15 (SPSS Inc., Chicago, IL) was used. The associations could be linear, logarithmic, potential, exponential, or polynomial. All functions were considered, and the most relevant one was retained. The mathematic function of the relation was individually determined through the least-squares fit and graphically represented (Fig. 3). In 166 of the children, it was quadratic and parabolic, which corresponds with a second-degree polynomial equation. Once the equation was defined, the optimal grip span was calculated as  $x \mid f'(x) = 0$ , where  $x$  equals the optimal grip

TABLE 1: Optimal Grip Span Determined in Boys (n = 123) for Each Hand Span\*

Hand Span (cm)	Optimal Grip Span for Right Hand (cm)	Optimal Grip Span for Left Hand (cm)	Optimal Grip Span† (cm)	p Value‡
15	4.2 ± 0.6	4.3 ± 0.5	4.2	.958
16	4.1 ± 0.6	4.3 ± 0.6	4.2	.284
17	5.0 ± 0.4	4.9 ± 0.5	5	.906
18	5.1 ± 0.6	5.1 ± 0.4	5.1	.885
19	5.2 ± 0.5	5.3 ± 0.4	5.2	.973
20	5.3 ± 0.4	5.3 ± 0.4	5.3	.657

\*The precision of the hand span measure was 0.5 cm, and the value was rounded to the nearest whole centimeter.

†Optimal grip span obtained from the mean of right- and left-hand optimal grip span.

‡Comparison between optimal grip span obtained with right hand versus optimal grip span obtained with left hand for each hand span.

span (cm) and  $f(x)$  equals the hand grip strength (kg). In graphic terms, this corresponded with the maximum of the curves (Fig. 3). For nonpolynomial equations ( $n = 12$ ), the optimal grip span was graphically determined, and this corresponded with one of the extreme grip spans used for that particular subject. For those children in whom there was no statistically significant association between hand grip strength and grip span ( $n = 15$ ; 6 boys and 9 girls aged 6 and 7 y), the average of the chosen grip spans was retained.

#### Determination of the optimal grip span for a given hand size:

By using the statistical software SPSS ver. 15, we studied whether optimal grip spans were significantly related to hand spans. In case of a significant relationship, the least-squares fit was used to calculate the mathematical function relating both variables. These equations allow the establishment of the optimal grip span for a given hand span. In case of a non-significant relationship, the conclusion is that optimal grip spans are not related to hand spans.

**Usefulness and reliability of the optimal grip span:** To confirm the usefulness of the optimal grip span when measuring hand grip strength in children aged 6 to 12 years, an additional group of 28 children (18 boys, 10 girls) of the same ages volunteered to perform the hand grip strength test at 3 grip spans: optimal grip span, 1 cm below the optimal grip span, and 1 cm above the optimal grip span. Each child performed the test following the same protocol as explained above. For each hand, the best result at each grip span was retained. To confirm the reliability of measurements of hand grip strength at the optimal grip span, 22 (17 boys, 5 girls) of the previous 28 children performed the test at the optimal grip span 2 hours later.

#### Statistical Analysis

The hand span, hand grip strength, and the optimal grip span obtained for each hand span and gender was compared by 1-way analysis of variance (ANOVA). Bivariate

correlation analysis was performed to examine the relationship between optimal grip span and hand span for each hand and gender. In case of an association, the mathematical function defining the association was calculated through the least-squares fit. For confirming the usefulness of measuring hand grip strength at the optimal grip span, 1 cm below the optimal grip span, and 1 cm above the optimal grip span, a 1-way ANOVA for repeated measures was used. The reliability coefficient of hand grip strength measured at the optimal grip span on 2 different occasions was calculated; values were compared through 1-way ANOVA for repeated measures and correlated through parametric bivariate correlation analysis. The  $\alpha$  error was fixed at .05.

#### RESULTS

The children completed all the tests satisfactorily. The measured hand span (mean  $\pm$  SD) was  $17.8 \pm 1.5$  cm for boys ( $n = 123$ ) and  $17.2 \pm 1.4$  cm for girls ( $n = 70$ ) ( $p = .004$ ). Boys obtained higher values of hand grip strength at each grip span than did girls (all  $p \leq .004$ ) (data not shown). In both genders, and for both hands, an optimal grip span was obtained. The optimal grip span for each hand span for boys and girls is presented in Tables 1 and 2, respectively. The optimal grip span was not significantly different between right and left hands ( $p > .06$ ); therefore, the mean value was retained and used for subsequent analysis.

The hand span and optimal grip span showed a significant linear association in the studied children ( $y = 0.2674x + 0.0883$  cm;  $r = 0.98$ ,  $p < .001$ ), where  $x$  is the hand span (maximal width between first and fifth fingers), and  $y$  is the optimal grip span at which the dynamometer should be adjusted before the test. The equation relating grip span as a function of hand span in boys is formulated as  $y = 0.2506x + 0.44$  cm ( $r = 0.97$ ,  $p = .002$ ). A simplification of this algorithm would be the following:  $y = x/4 + 0.44$  cm (Fig. 4). The equation relating grip span as a function of hand span in girls is formulated as  $y = 0.3x -$

TABLE 2: Optimal Grip Span Determined in Girls (n = 70) for Each Hand Span\*

Hand Span (cm)	Optimal Grip Span for Right Hand (cm)	Optimal Grip Span for Left Hand (cm)	Optimal Grip Span† (cm)	p Value‡
15	4.1 ± 0.5	4.2 ± 0.4	4.2	.549
16	4.3 ± 0.6	4.1 ± 0.4	4.2	.413
17	4.4 ± 0.4	4.3 ± 0.3	5	.759
18	5.3 ± 0.5	5.2 ± 0.5	5.1	.657
19	5.3 ± 0.5	5.1 ± 0.5	5.2	.492

\*The precision of the hand span measure was 0.5 cm, and the value was rounded to the nearest whole centimeter.

†Optimal grip span obtained from the mean of right- and left-hand optimal grip span.

‡Comparison between optimal grip span obtained with right hand versus optimal grip span obtained with left hand for each hand span.

0.52 cm ( $r = 0.96$ ,  $p = .008$ ) (Fig. 4). The optimal grip span for each hand span in boys and girls calculated from the equations provided is presented in Table 3.

The hand grip strength obtained at the optimal grip span was significantly higher than the strength obtained when the grip was set 1 cm below or 1 cm above the optimal grip span in both hands and genders ( $p < .02$ ; Fig. 5).

Twenty-two children (17 boys and 5 girls) from the previous 28 repeated the test 2 hours later at optimal grip span. The results showed a reliability coefficient of 0.972 and 0.985 for right and left hands, respectively. Moreover, the 1-way ANOVA for repeated measures did not show statistical difference between test and retest ( $p = .991$  and  $.402$  for right and left hands, respectively). A significant correlation between test and retest was obtained for right ( $r = 0.923$ ,  $p < .001$ ) and left ( $r = 0.971$ ,  $p < .001$ ) hands at the optimal grip span.

## DISCUSSION

This study shows that there is an optimal grip span at which the standard dynamometer should be adjusted when measuring hand grip strength in children. In both genders, the optimal grip span is influenced by hand span, which implies the need of adjusting the grip span of the dynamometer to the hand span. For that purpose,

gender-specific equations are proposed. This study shows how to mathematically adjust the grip span of the dynamometer to the hand span when measuring hand grip strength in both boys and girls as young as 6 to 12 years.

These findings are in agreement with our previous reports in teenagers<sup>12</sup> and in adults.<sup>13</sup> The optimal grip was influenced by hand span in male and female teenagers as well as in adult women. The equation to calculate the optimal grip span from the measure of the hand span in male and female teenagers was  $y = x/7.2 + 3.1$  cm ( $r = 0.92$ ,  $p = .01$ ) and  $y = x/4 + 1.1$  cm ( $r = 0.93$ ,  $p = .02$ ), respectively.<sup>12</sup> In adult women, the equation was  $y = x/5 + 1.5$  cm ( $r = 0.97$ ,  $p = .004$ ), whereas in adult men, there was an optimal grip span for determining the maximum hand grip strength, but that optimal grip span was not hand span-dependent. Therefore, a fixed optimal grip span was proposed (5.5 cm).<sup>13</sup> Children have smaller hand spans and lower hand grip strength compared with teenagers and adults. The same applies to teenagers in comparison with adults. Because of these differences, one would expect that each age group may need a different optimal grip span when measuring hand grip strength. For the same reasons, gender-specific equations have also been proposed.




FIGURE 4: Association between hand span and optimal grip span in boys (n = 123) and girls (n = 70).

TABLE 3: Optimal Grip Span for Each Hand Span Calculated From the Equations Provided\*

Hand Size (cm)	Optimal Grip Span for Boys (cm)	Optimal Grip Span for Girls (cm)	Optimal Grip Span for Boys and Girls (cm)
15.0	4.2	4.0	4.1
15.5	4.3	4.1	4.2
16.0	4.4	4.3	4.4
16.5	4.6	4.4	4.5
17.0	4.7	4.6	4.6
17.5	4.8	4.7	4.8
18.0	5.0	4.9	4.9
18.5	5.1	5.0	5.0
19.0	5.2	5.2	5.2
19.5	5.3	5.3	5.3
20.0	5.5	5.5	5.4

\*For boys and girls,  $\gamma = 0.2674x + 0.0883$  cm ( $r = 0.98$ ,  $p < .001$ ); for boys,  $\gamma = x/4 + 0.44$  cm ( $r = 0.97$ ,  $p = .002$ ); for girls,  $\gamma = 0.3x - 0.52$  cm ( $r = 0.96$ ,  $p = .008$ ); where  $x$  is the hand span (maximal width between thumb and little finger, with 0.5-cm precision), and  $\gamma$  is the optimal grip span in centimeters.

There is a general agreement suggesting that both grip and hand spans affect maximum hand grip strength.<sup>8,10-19</sup> Oh and Radwin<sup>18</sup> reported that hand span affected maximal and submaximal hand grip strength. They found that hand span affects grip strength, grip force, and exertion level. Firrell and Crain<sup>19</sup> studied which setting of the dynamometer produced maximal grip strength and correlated that setting with characteristics of the individual. They reported that the majority of the hands (89%) had a maximal strength at setting II (of V) of a hand dynamometer, whereas no clear significant correlation between hand size and maximal setting was found. More recently, Ratamess et al<sup>17</sup> examined the strength performance of 6 common resistance-training exercises using

free-weight bars of 3 different thicknesses. They found significant differences in the strength performance between bars and concluded that the observed differences might be due to greater reliance on maximal grip strength and larger hand size.

Hand grip strength is currently used worldwide because it is a relatively cheap test that gives practical information on muscle, nerve, bone, or joint disorders.<sup>20-25</sup> In adults, hand grip strength has been proposed as a possible predictor of mortality and the expectancy of being able to live independently.<sup>26,27</sup> Moreover, hand grip strength is associated with bone mineral density,<sup>28-30</sup> impaired cognition,<sup>31</sup> nutritional status,<sup>32,33</sup> and cardiovascular disease risk factors.<sup>34,35</sup> Therefore, from a public health perspective,




FIGURE 5: Hand grip strength measured in right and left hands at optimal grip span, 1 cm below, and 1 cm above in boys ( $n = 18$ ) and girls ( $n = 10$ ) (age range, 6 to 12 y). Values are mean  $\pm$  standard error of the mean. \* $p \leq .01$  compared with 1 cm below and 1 cm above. † $p < .02$  compared with 1 cm below and 1 cm above.

it is important to standardize the procedure and increase the reliability, as otherwise the measurement error may be too large to detect actual changes in strength. It must be borne in mind that different kinds of dynamometers and postures might change the results. We do not know whether these findings can be directly transferred to measurements with other dynamometers.

## REFERENCES

- Ruiz JR, Ortega FB, Gutierrez A, Meusel D, Sjöström M, Castillo MJ. Health-related fitness assessment in childhood and adolescence; a European approach based on the AVENA, EYHS and HELENA studies. *J Public Health* 2006;14:269–277.
- EUROFIT. Committee of Experts on Sports Research. Hand-book for the EUROFIT tests of physical fitness. Strasburg, GE: Council of Europe, 1993:19–37.
- Ahrens W, Bammann K, de Henauw S, Halford J, Palou A, Pigeot I, et al. Understanding and preventing childhood obesity and related disorders—IDEFICS: a European multilevel epidemiological approach. *Nutr Metab Cardiovasc Dis* 2006;16:302–308.
- Richards LG, Olson B, Palmiter-Thomas P. How forearm position affects grip strength. *Am J Occup Ther* 1996;50:133–138.
- Su CY, Lin JH, Chien TH, Cheng KF, Sung YT. Grip strength in different positions of elbow and shoulder. *Arch Phys Med Rehabil* 1994;75:812–815.
- Mathiowetz V, Rennells C, Donahoe L. Effect of elbow position on grip and key pinch strength. *J Hand Surg* 1985;10A:694–697.
- Kraft GH, Detels PE. Position of function of the wrist. *Arch Phys Med Rehabil* 1972;53:272–275.
- Watanabe T, Owashi K, Kanauchi Y, Mura N, Takahara M, Ogino T. The short-term reliability of grip strength measurement and the effects of posture and grip span. *J Hand Surg* 2005;30A:603–609.
- Kato T, Miyamoto K, Shimizu K. Postural reaction during maximum grasping maneuvers using a hand dynamometer in healthy subjects. *Gait Posture* 2004;20:189–195.
- Fransson C, Winkel J. Hand strength: the influence of grip span and grip type. *Ergonomics* 1991;34:881–892.
- Härkönen R, Piirtomaa M, Alaranta H. Grip strength and hand position of the dynamometer in 204 Finnish adults. *J Hand Surg* 1993;18B:129–132.
- Ruiz JR, Espana-Romero V, Ortega FB, Sjoström M, Castillo MJ, Gutierrez A. Hand span influences optimal grip span in male and female teenagers. *J Hand Surg* 2006;31A:1367–1372.
- Ruiz-Ruiz J, Mesa JL, Gutierrez A, Castillo MJ. Hand size influences optimal grip span in women but not in men. *J Hand Surg* 2002;27A:897–901.
- Clerke AM, Clerke JP, Adams RD. Effects of hand shape on maximal isometric grip strength and its reliability in teenagers. *J Hand Ther* 2005;18:19–29.
- Boadella JM, Kuijjer PP, Sluiter JK, Frings-Dresen MH. Effect of self-selected handgrip position on maximal handgrip strength. *Arch Phys Med Rehabil* 2005;86:328–331.
- Blackwell JR, Kornatz KW, Heath EM. Effect of grip span on maximal grip force and fatigue of flexor digitorum superficialis. *Appl Ergon* 1999;30:401–405.
- Ratamess NA, Faigenbaum AD, Mangine GT, Hoffman JR, Kang J. Acute muscular strength assessment using free weight bars of different thickness. *J Strength Cond Res* 2007;21:240–244.
- Oh S, Radwin RG. Pistol grip power tool handle and trigger size effects on grip exertions and operator preference. *Hum Factors* 1993;35:551–569.
- Firrell JC, Crain GM. Which setting of the dynamometer provides maximal grip strength? *J Hand Surg* 1996;21A:397–401.
- Louhevaara V, Smolander J, Aminoff T, Korhonen O, Shen N. Cardiorespiratory responses to fatiguing dynamic and isometric hand-grip exercise. *Eur J Appl Physiol* 2000;82:340–344.
- Alperovitch-Najenson D, Carmeli E, Coleman R, Ring H. Handgrip strength as a diagnostic tool in work-related upper extremity musculoskeletal disorders in women. *ScientificWorld Journal* 2004;4:111–117.
- Schreuders TA, Roebroek M, van der Kar TJ, Soeters JN, Hovius SE, Stam HJ. Strength of the intrinsic muscles of the hand measured with a hand-held dynamometer: reliability in patients with ulnar and median nerve paralysis. *J Hand Surg* 2000;25B:560–565.
- Merkies IS, Schmitz PI, van der Meche FG, Samijn JP, van Doorn PA. Connecting impairment, disability, and handicap in immune mediated polyneuropathies. *J Neurol Neurosurg Psychiatry* 2003;74:99–104.
- Adams J, Burridge J, Mullee M, Hammond A, Cooper C. Correlation between upper limb functional ability and structural hand impairment in an early rheumatoid population. *Clin Rehabil* 2004;18:405–413.
- Wessel J, Kaup C, Fan J, Ehalt R, Ellsworth J, Speer C, et al. Isometric strength measurements in children with arthritis: reliability and relation to function. *Arthritis Care Res* 1999;12:238–246.
- Metter EJ, Talbot LA, Schrage M, Conwit R. Skeletal muscle strength as a predictor of all-cause mortality in healthy men. *J Gerontol A Biol Sci Med Sci* 2002;57:B359–365.
- Seguin R, Nelson ME. The benefits of strength training for older adults. *Am J Prev Med* 2003;25:141–149.
- Ozgoçmen S, Karaoglan B, Cimen OB, Yorgancıoğlu ZR. Relation between grip strength and hand bone mineral density in healthy women aged 30–70. *Singapore Med J* 2000;41:268–270.
- Di Monaco M, Di Monaco R, Manca M, Cavanna A. Handgrip strength is an independent predictor of distal radius bone mineral density in postmenopausal women. *Clin Rheumatol* 2000;19:473–476.
- Dixon WG, Lunt M, Pye SR, Reeve J, Felsenberg D, Silman AJ, et al. Low grip strength is associated with bone

- mineral density and vertebral fracture in women. *Rheumatology (Oxford)* 2005;44:642–646.
31. Raji MA, Kuo YF, Snih SA, Markides KS, Peek MK, Ottenbacher KJ. Cognitive status, muscle strength, and subsequent disability in older Mexican Americans. *J Am Geriatr Soc* 2005;53:1462–1468.
32. Chilima DM, Ismail SJ. Nutrition and handgrip strength of older adults in rural Malawi. *Public Health Nutr* 2001; 4:11–17.
33. Pieterse S, Manandhar M, Ismail S. The association between nutritional status and handgrip strength in older Rwandan refugees. *Eur J Clin Nutr* 2002;56:933–939.
34. Garcia-Artero E, Ortega FB, Ruiz JR, Mesa JL, Delgado M, Gonzalez-Gross M, et al. [Lipid and metabolic profiles in adolescents are affected more by physical fitness than physical activity (AVENA study)]. *Rev Esp Cardiol* 2007;60:581–588.
35. Ozdirenc M, Biberoglu S, Ozcan A. Evaluation of physical fitness in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract* 2003;60:171–176.